

Crossbow*

Specialty Herbicide

FOLIAR RATE CARD

Sprayer Size	1% Solution	1½% Solution	4% Solution ¹
1 gal.	1⅓ fl. oz.	2 fl. oz.	5⅓ fl. oz.
3 gal.	4 fl. oz.	6 fl. oz.	1 pt.
5 gal.	6⅔ fl. oz.	10 fl. oz.	1⅔ pt.
50 gal.	2 qt.	3 qt.	2 gal.
100 gal.	1 gal.	1½ gal.	4 gal.

1% SOLUTION CONTROLS:

Weeds – Lambsquarters, Common Ragweed, Wild Mustard, Annual Fleabane, Marestalk, Sunflowers, Burdock, Tansy Ragwort, Cocklebur, Broadleaf Plantain.

Brush – Blackberry, Honeysuckle, Multiflora Rose², Sumac, Willow.

1½% SOLUTION CONTROLS:

Weeds – Pokeweed, Ironweed, Musk Thistle, Bull Thistle, Hemp Dogbane, Vetch, Hemp Sesbania, Bedstraw, Clover, Buttercup, Canada Thistle, Dandelions, Docks, Field Pennycress, Ground Ivy, Marshelder, Mouse-Ear Chickweed, Oxalis, Redroot Pigweed, Wild Carrot, Wild Violets.

Suppresses: Goldenrod, Kochia, Milkweed and horsenettle.

Brush – Cherry (except black), Sweetgum, Ash, Alder, Hazel, Russian Olive, Elm, Elderberry, Aspen, Birch, Red Maples, Oaks (White and tan), Poison Oak, Poison Ivy, Scotch Broom, Tamarack, Sassafras, Willow, Black Locust and Ceanothus.

¹Primarily for basal uses – see label.

²NOTE: For best results with Multiflora Rose, apply 1% solution at early to mid-flowering stage. Outside that period, apply 1.5% solution retreatment may be necessary the following year.

Application of Crossbow* Herbicide

Crossbow can be used in almost any type of spray equipment, from one-gallon hand-held sprayers to professional 1000-gallon or more, custom sprayers. Basal applications can be used year round.

- **Timing:** For best results for foliar applications, apply when weeds and brush are actively growing. This is most likely to occur in the spring to early summer, when moisture and temperature are favorable.
- **While Spraying:** Keep up moderate agitation to maintain the emulsion.
- **After Spraying:** One year after treatment, check the treated area from time to time and spot treat missed plants and new seedlings that could reinfest the area.
- **Use Precaution:** Do not apply directly to, or otherwise permit it to come into direct contact with cotton, grapes, tobacco, vegetables, flowers, fruit or ornamental trees, or other desirable broadleaf plants and do not permit spray mists containing it to drift onto them.

Application should be made only when hazards from spray drift are at a minimum. Do not spray when wind is blowing toward susceptible crops or ornamental plants near enough to be injured.

NOTE: Regardless of the herbicide you use, READ AND FOLLOW ALL LABEL DIRECTIONS CAREFULLY.